

Please write clearly in	block capitals.	
Centre number	Candidate number	
Surname		
Forename(s)		
Candidate signature	I declare this is my own work.	/

GCSE CHEMISTRY

Higher Tier Paper 2

Time allowed: 1 hour 45 minutes

Materials

For this paper you must have:

- a ruler
- a scientific calculator
- the periodic table (enclosed).

Instructions

- Use black ink or black ball-point pen.
- Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.
- In all calculations, show clearly how you work out your answer.


Information

- The maximum mark for this paper is 100.
- The marks for questions are shown in brackets.
- You are expected to use a calculator where appropriate.
- You are reminded of the need for good English and clear presentation in your answers.



For Examiner's Use		
Question	Mark	
1		
2		
3		
4		
5		
6		
7		
8		
TOTAL		

IB/M/Jun22/E9






	Copper can be obtained by recycling scrap copper wire.	Do not write outside the box
01.2	Suggest why poly(butene) insulation must be removed from scrap copper wire before the copper is recycled. [1 mark]	
01.3	Describe how scrap copper wire can be recycled to make new copper water pipes. [2 marks]	
0 1.4	Suggest two reasons why recycling scrap copper is more sustainable than extracting copper from copper ores. [2 marks]	
	2	
	Question 1 continues on the next page	



	Copper sulfate is a compound of copper.	Do not write outside the box
	Copper sulfate solution contains copper(II) ions and sulfate ions.	
0 1.5	A solution can be added to copper sulfate solution to show the presence of copper(II) ions.	
	Name the solution added.	
	Give the result of the test. [2 marks	5]
	Name of solution added	_
		_
	Result	_
		_
0 1.6	Describe one test to show the presence of sulfate ions in copper sulfate solution.	
	Give the result of the test. [2 marks]	51
	Test	_
		_
	Result	
		_ 11







This is the method used.

- 1. Add 2.0 g of hydrated cobalt chloride to an empty test tube.
- 2. Measure the mass of the test tube and contents.
- 3. Heat the test tube and contents gently for 30 seconds.
- 4. Allow the test tube and contents to cool.
- 5. Measure the mass of the test tube and contents.
- 6. Repeat steps 3 to 5 until the mass of the test tube and contents does not change.

Table 1 shows the results.

Table 1

Total heating time in seconds	Mass of test tube and contents in grams
0	26.5
30	26.2
60	25.9
90	25.6
120	25.6



Do not write outside the


box

6

02.1	Determine the mass of the empty test tube. [1 mark]	Do not write outside the box
	Mass of empty test tube = g	
02.2	Explain why the mass of the test tube and contents decreased. [2 marks]	
02.3	Suggest why the test tube and contents were heated until the mass did not change. [1 mark]	
	Question 2 continues on the next page	

	Energy is taken in from the surroundings when hydrated cobalt chloride is heated.	Do not write outside the box
02.4	When 238 g of hydrated cobalt chloride is heated until the mass does not change, 88.1 kJ of energy is taken in.	
	The student heated 2.00 g of hydrated cobalt chloride until the mass did not change.	
	Calculate the energy taken in during this reaction.	
	Give your answer to 3 significant figures. [3 marks]	
	Energy taken in (3 significant figures) = kJ	
02.5	What type of reaction takes place when hydrated cobalt chloride is heated? [1 mark]	
		8









This question is about life cycle assessments (LCAs).

0 3 . 1 Milk bottles can be made from glass or from a polymer.

 Table 2 shows information about milk bottles of equal volume.

Table 2

10

	Glass	Polymer
Raw materials	Limestone Sand Sodium carbonate	Crude oil
Energy needed to process raw materials in kilojoules	6750	1710
Energy needed to manufacture bottle in kilojoules	750	90
Mass of bottle in grams	200	20
Mean number of times used during lifetime of bottle	25	1
One disposal method at end of useful life	Recycled to make different glass products	Recycled to make different polymer products

Evaluate the use of glass for milk bottles compared with the use of a polymer for milk bottles.

Use features of life cycle assessments (LCAs) in your answer.

Use Table 2.

[6 marks]

Do not write outside the



		Do not write outside the box
		JOX
0 3.2	Milk is also sold in cardboard cartons.	
	A carton is made using 40 cm ³ of cardboard.	
	The density of the cardboard is 0.40 g/cm ³ .	
	Calculate the mass of the carton.	
	Use the equation:	
	density = $\frac{1}{\text{volume}}$	
	[3 marks]	
	Mass = g	9





This question is about the fractions obtained from crude oil.

Crude oil is separated into fractions by fractional distillation.

The fractions obtained from crude oil include:

- Iubricating oil
- naphtha
- petroleum gases.

Table 3 shows the boiling point range of these fractions.

Table 3

Fraction	Boiling point range in °C
Lubricating oil	300–350
Naphtha	90–200
Petroleum gases	< 25

Explain how these fractions are obtained from crude oil by fractional distillation.

[4 marks]



04.2	Fractions from crude oil can be processed to produce feedstock for the petrochemical industry.	Do not write outside the box
	Which two are useful materials produced from this feedstock? [2 marks]	
	Tick (\checkmark) two boxes.	
	Alloys	
	Ceramics	
	Detergents	
	Fertilisers	
	Solvents	
	Another fraction obtained from crude oil is petrol.	
04.3	Petrol contains a hydrocarbon with the formula C_9H_{20}	
	Complete the equation for the complete combustion of C_9H_{20}	
	You should balance the equation. [2 marks]	
	C_9H_{20} + → +	
04.4	Petrol obtained from crude oil contains sulfur impurities. Explain why sulfur impurities are removed before petrol is burned in car engines. [2 marks]	



IB/M/Jun22/8462/2H

Fraction	Range of number of carbon atoms in each molecule
Kerosene	11–15
Heavy fuel oil	20–40

Table 4

A student predicted that heavy fuel oil is more viscous than kerosene.

The student's prediction was correct.

Justify the student's prediction.


[2 marks]

Do not write outside the



	The heavy fuel oil fraction can be processed to produce smaller hydrocarbon molecules.	Do not write outside the box
04.6	Name the process which produces smaller hydrocarbon molecules from heavy fuel oil. Give the conditions used in this process. [3 marks]	
	Name of process	
	Conditions	
0 4.7	Hydrocarbon molecules containing seven and eight carbon atoms can be produced when heavy fuel oil is processed.	
	Which pair of hydrocarbon molecules would both turn bromine water colourless? [1 mark]	
	Tick (\checkmark) one box.	
	C ₇ H ₁₄ and C ₈ H ₁₆	
	C ₇ H ₁₄ and C ₈ H ₁₈	
	C ₇ H ₁₆ and C ₈ H ₁₆	
	C ₇ H ₁₆ and C ₈ H ₁₈	16







0 5	This question is about water.	Do not write outside the box
0 5.1	Sewage is waste water.	
	Sewage contains organic matter.	
	Describe how sewage is treated to remove organic matter. [4 marks]	
	Question 5 continues on the next page	
	True area b	
	Turn over ►	



 Table 5 shows information about the composition and treatment of sea water and of ground water.

 Table 5

Sea water and ground water are treated to make them potable.

	Sea water	Ground water
Concentration of sodium ions and chloride ions before Process 1	Na⁺ : 0.5 mol/dm³ Cl⁻ : 0.5 mol/dm³	Na⁺ : 0.001 mol/dm³ Cl⁻ : 0.001 mol/dm³
Process 1	Reverse osmosis	Filtration
Concentration of sodium ions and chloride ions after Process 1	x	Na⁺ : 0.001 mol/dm³ Cl⁻ : 0.001 mol/dm³
Process 2	Add ozone	Expose to ultraviolet light

0 5 2 Sea water is desalinated during **Process 1**.

Which pair of concentrations could represent X in Table 5?

Tick (✓) one box.

Na ⁺ : 0.003 mol/dm ³	Cl ⁻ : 0.003 mol/dm ³	
Na ⁺ : 0.003 mol/dm ³	Cl⁻ : 0.5 mol/dm³	
Na ⁺ : 0.5 mol/dm ³	Cl ⁻ : 0.003 mol/dm ³	
Na ⁺ : 0.5 mol/dm ³	Cl⁻ : 0.5 mol/dm³	

Explain why the concentrations of sodium ions and of chloride ions in the ground water in **Table 5** are unchanged by **Process 1**.


IB/M/Jun22/8462/2H

Do not write outside the box

[1 mark]

0 5.4	Explain why the ground water in Table 5 requires Process 2 before the water is safe to drink. [2 marks]	Do not write outside the box
0 5.5	After treatment the ground water in Table 5 is sold by a company as pure water.	
	The ground water in Table 5 is not chemically pure because the water contains sodium ions and chloride ions.	
	Suggest what the company means by 'pure'.	
	[1 mark]	
0 5.6	Chlorine is also used to treat some ground water.	
	Describe the test for chlorine gas.	
	Give the result of the test.	
	[2 marks]	
	Test	
	Result	
		12








06.2	Explain the change in the percentage of gas in the region labelled B on Figure 3 . [2 marks]	Do not write outside the box
06.3	Compare the changes in the percentages of gases in the region labelled C on Figure 3 . [2 marks]	
06.4	What process caused the changes in the percentages of gases in the region labelled C on Figure 3 ? [1 mark]	
06.5	Natural gas is a fossil fuel. Describe how deposits of natural gas were formed.	
	[3 marks]	
		10



0 7	Ammonia is produced in the Haber process.	Do not write outside the box
	The raw materials for the Haber process are nitrogen and hydrogen.	
	The equation for the reaction is:	
	$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$	
0 7.1	Give the sources of the nitrogen and of the hydrogen used in the Haber process. [2 marks]	
	Nitrogen	
	Hydrogen	
0 7 . 2	How does the equation for the reaction show that the atom economy of the forward reaction is 100%?	
	[1 mark]	







Do not write outside the box


The Haber process uses a temperature of 450 °C and a pressure of 200 atmospheres.

Table 6 shows the percentage yield of ammonia produced at 450 °C using different pressures.

Pressure in atmospheres	Percentage (%) yield of ammonia
60	9
120	18
180	25
240	31
300	36
360	40
420	43

Table 6








20	
07 . 6 The equation for the production of ammonia in the Haber process	is:
$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$	
The forward reaction is exothermic.	
The conditions used are:	
 a temperature of 450 °C 	
 a pressure of 200 atmospheres 	
 the presence of an iron catalyst. 	
Explain why these conditions are chosen for economical production Haber process.	on of ammonia in the
You should include references to the rate of reaction and the posi	tion of equilibrium. [6 marks]

Do not write outside the box








IB/M/Jun22/8462/2H

08	This question is about the reaction between sodium thiosulfate solution and hydrochloric acid.		Do not write outside the box
	When hydrochloric acid is added to sodium thiosulfate solution, the mixture grabecomes cloudy.	adually	
	The equation for the reaction is:		
	$Na_2S_2O_3(aq)$ + 2 HCl(aq) \rightarrow 2 NaCl(aq) + H ₂ O(I) + SO ₂ (g)	S(s)	
08.1	Sulfur is produced in the reaction.		
	Why does the mixture become cloudy?	1 mark]	

A student investigated the effect of changing the concentration of sodium thiosulfate solution on the rate of the reaction.

Figure 6 shows the apparatus used.

Figure 6



A smaller percentage of light from the light source reaches the light sensor as the mixture becomes more cloudy.

This is the method used.


- 1. Measure 50 cm³ of 0.10 mol/dm³ sodium thiosulfate solution into the beaker.
- 2. Add 10 cm³ of hydrochloric acid to the sodium thiosulfate solution.
- 3. Immediately start a timer.
- 4. Record the percentage of light from the light source that reaches the light sensor every 20 seconds for 120 seconds.
- 5. Repeat steps 1 to 4 using 0.20 mol/dm³ sodium thiosulfate solution.

Question 8 continues on the next page



Turn over ►


Do not write outside the





0 8.2	The percentage of light reaching the light sensor decreases by 1% when 7.1 × 10^{-5} moles of sulfur is produced.	Do not w outside box
	Determine the rate of reaction in mol/s for the production of sulfur at 30 seconds.	
	You should draw a tangent on Figure 7 . [5 marks]	1
		-
		-
		-
		-
		-
		-
		-
		-
		-
	Rate = mol/s	6
	Eveloin why the acts of recetion channes hot year 0 and 00 accords	
0 8 . 3	Explain why the rate of reaction changes between 0 and 60 seconds.	
	Answer in terms of concentration.	
	Use Figure 7. [2 marks]	1
		_
		_
		-









		Do not write outside the
	The same student did the investigation again the next day.	box
	The student found that the same method produced different results for the percentage of light reaching the light sensor.	
08.6	How could the student improve the method so that the same percentages of light reached the light sensor? [1 mark]	
	Tick (✓) one box.	
	Record the percentage of light every 10 seconds.	
	Stop light from other sources reaching the light sensor.	
	Use a larger volume of sodium thiosulfate solution.	
	Use a more sensitive light sensor.	
08.7	The student improved the method so that similar results were obtained on different days.	
	What name is given to similar results obtained on different days under the same conditions by the same student?	
	[1 mark] Tick (✓) one box.	
	Anomalous	
	Precise	
	Repeatable	
	Reproducible	








Do not write outside the









Question number	Additional page, if required. Write the question numbers in the left-hand margin.




Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.





Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2022 AQA and its licensors. All rights reserved.





IB/M/Jun22/8462/2H